



NEW COMMON FIXED POINT RESULTS FOR DIGITAL TYPE EXPANSIVE MAPPINGS IN  
DIGITAL METRIC SPACE

DEEPAK KUMAR TIWARI<sup>1</sup>, SANGEETA PANDEY<sup>2</sup>, KHUSHBOO KUSHWAHA<sup>3</sup>

<sup>1</sup>Shri Sai Baba Aadarsh Mahavidyalaya, Ambikapur, Surguja (C.G.) India

<sup>2</sup>Rajeev Gandhi Govt. P.G. College, Ambikapur, Surguja (C.G.) India

<sup>3</sup>Shri Sai Baba Aadarsh Mahavidyalaya, Ambikapur, Surguja (C.G.) India

DOI:[10.33329/bomsr.11.2.57](https://doi.org/10.33329/bomsr.11.2.57)



**ABSTRACT**

Information Technology plays a vital role in every step of human activity in the 21<sup>st</sup> century. In this era, we can't imagine our life without technology because the virtual communication takes place over traditional communication by using various ways like video conferencing, telephoning, Emailing, Image processing, etc. On the other hand fixed point theory is a very traditional topic of pure part of Mathematics which plays a vital role in many disciplinary of Mathematics such as Topology, Physics, Chemistry, Biology, Computer Science, etc. In the second half years of the 20<sup>th</sup> century, researchers tried to provide a digital version of this theory. A digital image is the form of pixels and we can consider the distance between pixels by a distance function. As the distance between pixels is very least then we find a good image quality while the distance is maximum then we obtain a poor quality image. In this paper, we established existence of some new fixed point results by taking two self-mappings for expansive mappings in digital metric space.

**Keywords:** Fixed Point, Digital image, Digital type expansive mappings

**Introduction**

The fixed point theory (FPT) is now a promising area in mathematics especially in nonlinear

functional analysis because it has wide applicability in various fields of pure and applied mathematics as well as in other fields like Physical science, Life science Economics, etc. The field of the FPT is expanding its domain, thereby leading to the emergence of a plethora of techniques and ideas. The fixed point theory is one of the very effective and fruitful tools in Mathematics which has huge applications inside as well as outside mathematics. It is an influential field of specialization and has grown into a full branch of Mathematics within more than a hundred years out of multitude problems occurring in diversified fields. It is very difficult to imagine its applications in many fields.

Nowadays the human world has become completely digital. In every step of life humans use technology in every sector such as Medical, Education, Defence, etc. From these Image processing is a promising area in which FPT is more applicable.

Fixed Point theory for digital image firstly studied by A. Rosenfield [10]. In 2017, Jyoti [6] proved a fixed point theorem for expansive mapping on complete digital metric space. In recent years, many authors discussed on fixed point result in view of digital images and satisfy certain digital type contractive and expansive conditions with applications [1,3,5,6,7,8,9,11,12]. In this paper we establish new fixed point results for two self mappings satisfy digital type expansive condition in digital metric space. I hope this result extends many well known results. But, before proving the main result we need to explain some prerequisites.

## 2. PREREQUISITES:


**Definition 2.1.[1]:** Suppose  $Z^n$  the set of all integers defined by  $Z^n = \{(x_1, x_2, x_3, \dots, x_n) : x_i \in Z, 1 \leq i \leq n\}$  and called as the set of all lattice point in the  $n$ -dimensional Euclidean space. Suppose  $X \subseteq Z^n \forall n \in Z$  and  $\alpha$  represent and adjacency relation for the members of  $X$  then  $(X, \alpha)$  is called **digital image**.

**Definition 2.2.[1]:** Assume  $p, q \in Z^n$  Where  $p = (p_1, p_2, p_3 \dots p_n)$  and  $q = (q_1, q_2, q_3 \dots q_n)$  Let  $n \in Z$  such that  $1 \leq \alpha \leq n$  then we say  $p$  and  $q$  are  $\alpha$ -adjacent in  $Z^n$  if there exist at most  $\alpha$  indices  $i$  such that  $|p_i - q_i| \leq 1$  and for all other indices  $j$  such that  $|p_j - q_j| \neq 1$  we obtain  $p_j - q_j$ . From this definition following statement can be obtained

For given  $p \in Z^n$  the number of points  $q \in Z^n$  which are  $\alpha$ -adjacent to  $p$  is denoted by  $K(\alpha, n)$  Where  $K(\alpha, n)$  is independent to  $p$ .


For our convenience we write  $K = K(\alpha, n)$ , then we have following cases

- 1) If  $n = 1$  then assume  $p \in Z$  then  $\alpha$  will have only one value  $\alpha = 1$ . Then in this case  $K = K(1, 1) = 2$ . So in this case  $p - 1$  and  $p + 1$  only two points are 1- adjacent to  $p \in Z$  also  $q$  is 1-adjacent to  $p$  iff  $|p - q| = 1$
- 2) Let  $n = 2$  then assume  $p \in Z^{2++}$  so  $\alpha$  will have  $\alpha = 1, 2$  if  $\alpha = 2$  then 2 adjacent to  $p = (p_1, p_2)$ . In this case  $(p_1 \pm 1, p_2) (p_1, p_2 \pm 1) (p_1 \pm 1, p_2 \pm 1) (p_1 + 1, p_2 \pm 1)$  are only points of 2- adjacent to  $p$  such that  $K = K(2, 2) = 8$



**1-adjacency**

**4-adjacency**



**8-adjacency**

Similarly  $n = 3$  then.  $K = K(3,3) = 26$ . In general for study  $n$ -D digital image. If  $1 \leq \alpha \leq n$  then  $K = K(\alpha, n)$  is given by

$$K(\alpha, n) = \sum_{i=n-\alpha}^{n-1} 2^{n-i} C_i^n \qquad C_i^n = \frac{n!}{(n-i)!i!}$$

**Definition 2.3. [1]:** Let  $(X, k) \subseteq Z^n$  be a digital image. Define a function  $d : X \times X \rightarrow [0, \infty)$  by,

$$d(p, q) = \left[ \sum_{i=1}^n (p_i - q_i)^2 \right]^{\frac{1}{2}}$$

Then we have the following properties satisfied by  $d$  for all  $x, y, z \in X$ .

- DMS<sub>1</sub>       $d(x, y) \geq 0$
- DMS<sub>2</sub>       $d(x, y) = 0 \Leftrightarrow x = y$
- DMS<sub>3</sub>       $d(x, y) = d(y, x)$
- DMS<sub>4</sub>       $d(x, y) \leq d(x, z) + d(z, y)$

The digital image  $(X, k)$  together with the function  $d$  is called a digital metric space with  $k$ -adjacency. It is denoted by  $(X, d, k)$ .

i.e. "**A metric space equipped with digital image is called digital metric space.**"

**Definition 2.4. [1]:** Let  $(X, d, k)$  be a digital metric space then sequence  $\{x_n\}$  of the points of  $(X, d, k)$  is called

- 2.4.1. Cauchy sequences** in  $X$  if  $\forall \epsilon > 0, \exists n_0 \in Z_+$  such that  $d(x_n, x_m) < \epsilon, \forall n, m > n_0$ .
- 2.4.2. converges** to a limit  $L$  in  $X$  if for given  $\epsilon > 0, \exists n_0 \in Z_+$  such that  $d(x_n, L) < \epsilon, \forall n, > n_0$ .
- 2.4.3. complete** if every Cauchy sequence  $\{x_n\}$  of point of  $(X, d, k)$  converges to a point  $L$  of  $(X, d, k)$ .

**Definition 2.5[1]:** Let  $(X, d, k)$  be a digital metric space and  $T : X \rightarrow X$  be a mapping then a point  $x \in X$  is called a digital fixed point of  $T$  if  $x$  is mapped into itself, i.e.  $T(x) = x$ .

**Theorem [6]:** Let  $T : (X, d, k) \rightarrow (X, d, k)$  be a mapping on a complete digital metric space  $X$ . Let  $T$  be onto and satisfy  $d(Tx, Ty) \geq \lambda d(x, y) \forall x \in X$  and  $\lambda > 1$ . Then  $T$  has a fixed point in  $X$ . In this condition  $T$  is known as expansive mapping

### 3. MAIN RESULTS

In this section we proof five theorems with different type digital expansive conditions

**Theorem 3.3.1:** Let  $(X, k)$  be a digital image where  $X \subseteq Z^n$  and  $k$  is an adjacency relation in  $X$ . Let  $(X, d, k)$  be a complete digital metric space and suppose  $T_1, T_2 : X \rightarrow X$  be any two continuous and onto mapping satisfying the condition

$$d(T_1x, T_2y) \geq \alpha d(x, y) + \beta [d(x, T_1x) + d(y, T_2y)]$$

for all  $x, y \in X$ , where  $\alpha > 0, \beta \in \left[ \frac{1}{2}, 1 \right]$  are constant, with  $\alpha + \beta > 1$ . Then  $T_1$  and  $T_2$  have a common fixed point in  $X$ .

**Proof:** Let  $x_0 \in X$  is arbitrary point. Since  $T_1$  and  $T_2$  be onto so there exist  $x_0 \in X$  and  $x_1 \in X$  such that  $T_1(x_1) = x_0, T_2(x_2) = x_1$ .

In this way, we define the sequences  $\{x_{2n}\}$  and  $\{x_{2n+1}\}$  by

$$x_{2n} = T_1x_{2n+1} \text{ for } n = 0, 1, 2, 3 \dots \dots \text{and}$$

$$x_{2n+1} = T_2x_{2n+2} \text{ for } n = 0, 1, 2, 3 \dots \dots$$

Now put  $x = x_{2n+1}$  and  $y = x_{2n+2}$ , we have

$$d(x_{2n}, x_{2n+1}) = d(T_1x_{2n+1}, T_2x_{2n+2})$$

$$\begin{aligned} \Rightarrow d(x_{2n}, x_{2n+1}) &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta [d(x_{2n+1}, T_1x_{2n+1}) + d(x_{2n+2}, T_2x_{2n+2})] \\ &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta [d(x_{2n+1}, x_{2n}) + d(x_{2n+2}, x_{2n+1})] \\ &= (\alpha + \beta) d(x_{2n+1}, x_{2n+2}) + \beta d(x_{2n+1}, x_{2n}) \end{aligned}$$

$$\Rightarrow (1 - \beta) d(x_{2n}, x_{2n+1}) \geq (\alpha + \beta) d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n}, x_{2n+1}) \geq \frac{\alpha + \beta}{1 - \beta} d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq \frac{1 - \beta}{\alpha + \beta} d(x_{2n}, x_{2n+1})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq h d(x_{2n}, x_{2n+1}), \text{ where } h = \frac{1 - \beta}{\alpha + \beta}, 0 \leq h < 1.$$

In general

$$d(x_{2n}, x_{2n+1}) \leq h d(x_{2n-1}, x_{2n}) \leq \dots \dots \leq h^{2n} d(x_0, x_1)$$

So, for  $n < m$  we have

$$\begin{aligned} d(x_{2n}, x_{2m}) &\leq d(x_{2n}, x_{2n+2}) + \dots \dots d(x_{2m-1}, x_{2m}) \\ &\leq (h^{2n} + h^{2n+1} + \dots \dots \dots + h^{2m-1}) d(x_0, x_1) \\ &= \frac{h^{2n}}{1 - h} d(x_0, x_1) \end{aligned}$$

Let  $c \geq 0$  be given, choose a natural number  $N$  such that  $\frac{h^{2n}}{1 - h} d(x_0, x_1) \leq c$ , for all  $n \geq N$ . Thus  $d(x_{2n}, x_{2m}) \leq c$ , for  $m > n$ . Therefore  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, d, k)$ .

Since  $(X, d, k)$  is a complete digital metric space, there exist  $z \in X$  such that  $x_{2n} \rightarrow z$  as  $n \rightarrow \infty$ . Also if  $T_1$  is continuous, then

$$d(T_1 z, z) \leq d(T_1 x_{2n+1}, T_1 z) + d((T_1 x_{2n+1}), z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since  $x_{2n} \rightarrow z$  and  $T_1 x_{2n+1} \rightarrow T_1 z \rightarrow z$  as  $n \rightarrow \infty$ . Therefore  $d(T_1 z, z) = 0$ . This implies that,  $T_1 z = z$ . Hence  $z$  is a fixed point of  $T_1$

Similarly, it can be established that  $T_2 z = z$ . Therefore,  $T_1 z = z = T_2 z$ . Thus  $z$  is the common fixed point of pair of maps  $T_1$  and  $T_2$ . This completes the proof.

**Theorem 3.2:** Let  $(X, k)$  be a digital image where  $X \subseteq Z^n$  and  $k$  is an adjacency relation in  $X$ . Let  $(X, d, k)$  be a complete digital metric space and suppose  $T_1, T_2 : X \rightarrow X$  be any two continuous and onto mapping satisfying the condition

$$d(T_1 x, T_2 y) \geq \alpha d(x, y) + \beta [d(x, T_2 y) + d(y, T_1 x)] \dots \dots \dots 3.1.2$$

for all  $x, y \in X$ , where  $\alpha > 0, \frac{1}{2} \leq \beta \leq 1$  are constant, with  $\alpha + \beta > 1$ . Then  $T_1$  and  $T_2$  have a common fixed point in  $X$ .

**Proof:** Let  $x_0$  be an arbitrary point in  $X$ . Since  $T_1$  and  $T_2$  be onto (surjective), there exist  $x_0 \in X$  and  $x_1 \in X$  such that

$$T_1(x_1) = x_0, T_2(x_2) = x_1$$

In this way, we define the sequences  $\{x_{2n}\}$  and  $\{x_{2n+1}\}$  by

$$x_{2n} = T_1 x_{2n+1} \text{ for } n = 0, 1, 2, 3 \dots \dots$$

$$\text{and } x_{2n+1} = T_2 x_{2n+2} \text{ for } n = 0, 1, 2, 3 \dots \dots$$

Note that, If  $x_{2n} = x_{2n+1}$  some  $n \geq 1$ , then it is fixed point of  $T_1$  and  $T_2$ .

Now putting  $x = x_{2n+1}$  and  $y = x_{2n+2}$ , we have

$$\begin{aligned} d(x_{2n}, x_{2n+1}) &= d(T_1 x_{2n+1}, T_2 x_{2n+2}) \\ d(x_{2n}, x_{2n+1}) &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta [d(x_{2n+1}, T_2 x_{2n+2}) + d(x_{2n+2}, T_1 x_{2n+1})] \\ &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta [d(x_{2n+1}, x_{2n+1}) + d(x_{2n+2}, x_{2n})] \\ &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta [d(x_{2n+2}, x_{2n+1}) + d(x_{2n+1}, x_{2n})] \\ &= (\alpha + \beta) d(x_{2n+1}, x_{2n+2}) + \beta d(x_{2n+1}, x_{2n}) \end{aligned}$$

$$\Rightarrow (1 - \beta) d(x_{2n}, x_{2n+1}) \geq (\alpha + \beta) d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n}, x_{2n+1}) \geq \frac{\alpha + \beta}{1 - \beta} d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq \frac{1 - \beta}{\alpha + \beta} d(x_{2n}, x_{2n+1})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq h d(x_{2n}, x_{2n+1}), \text{ where } h = \frac{1 - \beta}{\alpha + \beta}, 0 \leq h < 1.$$

In general

$$d(x_{2n}, x_{2n+1}) \leq h d(x_{2n-1}, x_{2n}) \leq \dots \dots \dots h^{2n} d(x_0, x_1)$$

So, for  $n < m$ , we have

$$\begin{aligned}
 d(x_{2n}, x_{2m}) &\leq d(x_{2n}, x_{2n+2}) + \dots + d(x_{2m-1}, x_{2m}) \\
 &\leq (h^{2n} + h^{2n+1} + \dots + h^{2m-1})d(x_0, x_1) \\
 &\leq \frac{h^{2n}}{1-h} d(x_0, x_1)
 \end{aligned}$$

Let  $0 \leq c$  be given, choose a natural number  $N_1$  such that  $\frac{h^{2n}}{1-h} d(x_0, x_1) \leq c$ , for all  $n \geq N_1$ . Thus  $d(x_{2n}, x_{2m}) \leq c$ , for  $n < m$ . Therefore  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, d)$ . Since  $(X, d)$  is a complete digital metric space, there exist  $z \in X$  such that  $x_{2n} \rightarrow z$  as  $n \rightarrow \infty$ . If  $T_1$  is continuous, then

$$d(T_1 z, z) \leq d(T_1 x_{2n+1}, T_1 z) + d((T_1 x_{2n+1}), z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since  $x_{2n} \rightarrow z$  and  $T_1 x_{2n+1} \rightarrow T_1 z$  as  $n \rightarrow \infty$ . Therefore  $d(T_1 z, z) = 0$ . This implies that,  $T_1 z = z$ . Hence  $z$  is a fixed point of  $T_1$ .

Similarly, it can be established that  $T_2 z = z$ . Therefore,  $T_1 z = z = T_2 z$ . Thus  $z$  is the common fixed point of pair of maps  $T_1$  and  $T_2$ . This completes the proof.

**Theorem 3.3:** Let  $(X, k)$  be a digital image where  $X \subseteq Z^n$  and  $k$  is an adjacency relation in  $X$ . Let  $(X, d, k)$  be a complete digital metric space and suppose  $T_1, T_2 : X \rightarrow X$  be any two continuous and onto mapping satisfying the condition

$$d(T_1 x, T_2 y) \geq \alpha d(x, y) + \beta d(x, T_1 x) + \gamma d(y, T_2 y) + \eta [d(x, T_1 x) + d(y, T_2 y)] \dots 3.3.1$$

for all  $x, y \in X$ , where  $\alpha \geq -1, \beta > 0, \gamma \leq \frac{1}{2}$  and  $\frac{1}{2} < \eta \leq 1$  are constant, with  $\alpha + \beta + \gamma + \eta > 1$ . Then  $T_1$  and  $T_2$  have a common fixed point in  $X$ .

**Proof:** Let  $x_0$  be an arbitrary point in  $X$ . Since  $T_1$  and  $T_2$  be onto (surjective), there exist  $x_1 \in X$  and  $x_2 \in X$  such that

$$T_1(x_1) = x_0, \quad T_2(x_2) = x_1$$

In this way, we define the sequences  $\{x_{2n}\}$  and  $\{x_{2n+1}\}$  by

$$x_{2n} = T_1 x_{2n+1} \text{ for } n = 0, 1, 2, 3 \dots$$

and 
$$x_{2n+1} = T_2 x_{2n+2} \text{ for } n = 0, 1, 2, 3 \dots$$

Now putting  $x = x_{2n+1}$  and  $y = x_{2n+2}$ .

Then we have,

$$\begin{aligned}
 d(x_{2n}, x_{2n+1}) &= d(T_1 x_{2n+1}, T_2 x_{2n+2}) \\
 \Rightarrow d(x_{2n}, x_{2n+1}) &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta d(x_{2n+1}, T_1 x_{2n+1}) + \gamma d(x_{2n+2}, T_2 x_{2n+2}) \\
 &\quad + \eta [d(x_{2n+1}, T_1 x_{2n+1}) + d(x_{2n+2}, T_2 x_{2n+2})] \\
 &\geq \alpha d(x_{2n+1}, x_{2n+2}) + \beta d(x_{2n+1}, x_{2n}) + \gamma d(x_{2n+2}, x_{2n+1}) \\
 &\quad + \eta [d(x_{2n+1}, x_{2n}) + d(x_{2n+2}, x_{2n+1})] \\
 &= (\alpha + \gamma + \eta) d(x_{2n+1}, x_{2n+2}) + (\beta + \eta) d(x_{2n+1}, x_{2n})
 \end{aligned}$$

$$\Rightarrow [1 - (\beta + \eta)] d(x_{2n}, x_{2n+1}) \geq (\alpha + \gamma + \eta) d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n}, x_{2n+1}) \geq \frac{\alpha + \gamma + \eta}{1 - (\beta + \eta)} d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq \frac{1-(\beta+\eta)}{\alpha+\gamma+\eta} d(x_{2n}, x_{2n+1})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq hd(x_{2n}, x_{2n+1}), \text{ where } h = \frac{1-(\beta+\eta)}{\alpha+\gamma+\eta}, 0 \leq h \leq 1.$$

In general

$$d(x_{2n}, x_{2n+1}) \leq hd(x_{2n-1}, x_{2n}) \leq \dots, \dots, \dots h^{2n}d(x_0, x_1)$$

So, for  $n < m$ , we have

$$\begin{aligned} d(x_{2n}, x_{2m}) &\leq d(x_{2n}, x_{2n+2}) + \dots, \dots, \dots d(x_{2m-1}, x_{2m}) \\ &\leq (h^{2n} + h^{2n+1} + \dots + h^{2m-1})d(x_0, x_1) \\ &\leq \frac{h^{2n}}{1-h}d(x_0, x_1) \end{aligned}$$

Let  $0 \leq c$  be given, choose a natural number  $N_1$  such that  $\frac{h^{2n}}{1-h}d(x_0, x_1) \leq c$ , for all  $n \geq N_1$ . Thus  $d(x_{2n}, x_{2m}) \leq c$ , for  $n < m$ . Therefore  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, d, k)$ . Since  $(X, d, k)$  is a complete digital metric space, there exist  $z \in X$  such that  $x_{2n} \rightarrow z$  as  $n \rightarrow \infty$ .

If  $T_1$  is continuous, then

$$d(T_1 z, z) \leq d(T_1 x_{2n+1}, T_1 z) + d((T_1 x_{2n+1}, z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since  $x_{2n} \rightarrow z$  and  $T_1 x_{2n+1} \rightarrow T_1 z$  as  $n \rightarrow \infty$ . Therefore  $d(T_1 z, z) = 0$ . This implies that,  $T_1 z = z$ . Hence  $z$  is a fixed point of  $T_1$ .

Similarly, it can be established that  $T_2 z = z$ . Therefore,  $T_1 z = z = T_2 z$ . Thus  $z$  is the common fixed point of pair of maps  $T_1$  and  $T_2$ . This completes the proof.

**Theorem 3.4:** Let  $(X, k)$  be a digital image where  $X \subseteq Z^n$  and  $k$  is an adjacency relation in  $X$ . Let  $(X, d, k)$  be a complete digital metric space and suppose  $T_1, T_2 : X \rightarrow X$  be any two continuous and onto mapping satisfying the condition

$$d(T_1 x, T_2 y) \geq \alpha[d(x, y) + d(x, T_1 x) + d(y, T_2 y)] + [\beta d(x, T_2 y) + d(y, T_1 x)] \dots 3.1.4$$

for all  $x, y \in X$ , where  $\alpha \geq 0, \beta < 1$  are constant, with  $\alpha + \beta > 1$ . Then  $T_1$  and  $T_2$  have a common fixed point in  $X$ .

**proof :** Let  $x_0$  be an arbitrary point in  $X$ . Since  $T_1$  and  $T_2$  be onto (surjective), there exist  $x_0 \in X$  and  $x_1 \in X$  such that

$$T_1(x_1) = x_0, T_2(x_2) = x_1$$

In this way, we define the sequences  $\{x_{2n}\}$  and  $\{x_{2n+1}\}$  by

$$x_{2n} = T_1 x_{2n+1} \text{ for } n = 0, 1, 2, 3, \dots$$

$$\text{and } x_{2n+1} = T_2 x_{2n+2} \text{ for } n = 0, 1, 2, 3, \dots$$

Note that, If  $x_{2n} = x_{2n+1}$  some  $n \geq 1$ , then it is fixed point of  $T_1$  and  $T_2$ .

Now putting  $x = x_{2n+1}$  and  $y = x_{2n+2}$ , we have

$$\begin{aligned} d(x_{2n}, x_{2n+1}) &= d(T_1 x_{2n+1}, T_2 x_{2n+2}) \\ &\geq \alpha[d(x_{2n+1}, x_{2n+2}) + d(x_{2n+1}, T_1 x_{2n+1}) + d(x_{2n+2}, T_2 x_{2n+2})] \end{aligned}$$

$$\begin{aligned}
 & + \beta[(x_{2n+1}, T_2x_{2n+2}) + (x_{2n+2}, T_1x_{2n+1})] \\
 \geq & \alpha[d(x_{2n+1}, x_{2n+2}) + d(x_{2n+1}, x_{2n}) + d(x_{2n+2}, x_{2n+1})] \\
 & + \beta[(x_{2n+1}, x_{2n+1}) + (x_{2n+2}, x_{2n})] \\
 \geq & \alpha[2d(x_{2n+1}, x_{2n+2}) + d(x_{2n+1}, x_{2n})] \\
 & + \beta[d(x_{2n+1}, x_{2n+2}) + d(x_{2n+2}, x_{2n+1}) + d(x_{2n+1}, x_{2n})] \\
 = & (2\alpha + 2\beta)d(x_{2n+1}, x_{2n+2}) + (\alpha + \beta)d(x_{2n}, x_{2n+1}) \\
 \Rightarrow & [1 - (\alpha + \beta)]d(x_{2n}, x_{2n+1}) \geq (2\alpha + 2\beta) d(x_{2n+1}, x_{2n+2}) \\
 \Rightarrow & d(x_{2n}, x_{2n+1}) \geq \frac{(2\alpha + 2\beta)}{1 - (\alpha + \beta)} d(x_{2n+1}, x_{2n+2}) \\
 \Rightarrow & d(x_{2n+1}, x_{2n+2}) \leq \frac{1 - (\alpha + \beta)}{(2\alpha + 2\beta)} d(x_{2n}, x_{2n+1})
 \end{aligned}$$

Therefore,  $d(x_{2n+1}, x_{2n+2}) \leq hd(x_{2n}, x_{2n+1})$ , where  $h = \frac{1 - (\alpha + \beta)}{(2\alpha + 2\beta)}$ ,  $0 \leq h \leq 1$ .

In general

$$d(x_{2n}, x_{2n+1}) \leq hd(x_{2n-1}, x_{2n}) \leq \dots \dots h^{2n}d(x_0, x_1)$$

So for  $n < m$ , we have

$$\begin{aligned}
 d(x_{2n}, x_{2m}) & \leq d(x_{2n}, x_{2n+2}) + \dots \dots + d(x_{2m-1}, x_{2m}) \\
 & \leq (h^{2n} + h^{2n+1} + \dots \dots + h^{2m-1})d(x_0, x_1) \\
 & \leq \frac{h^{2n}}{1-h} d(x_0, x_1)
 \end{aligned}$$

Let  $0 \leq c$  be given, choose a natural number  $N_1$  such that  $\frac{h^{2n}}{1-h}d(x_0, x_1) \leq c$ , for all  $n \geq N_1$ . Thus  $d(x_{2n}, x_{2m}) \leq c$ , for  $n < m$ . Therefore  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, d, k)$ . Since  $(X, d, k)$  is a complete digital metric space, there exist  $z \in X$  such that  $x_{2n} \rightarrow z$  as  $n \rightarrow \infty$ . If  $T_1$  is continuous, then

$$d(T_1 z, z) \leq d(T_1 x_{2n+1}, T_1 z) + d((T_1 x_{2n+1}), z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since  $x_{2n} \rightarrow z$  and  $T_1 x_{2n+1} \rightarrow T_1 z$  as  $n \rightarrow \infty$ . Therefore  $d(T_1 z, z) = 0$ . This implies that,  $T_1 z = z$ . Hence  $z$  is a fixed point of  $T_1$ .

Similarly, it can be established that  $T_2 z = z$ . Therefore,  $T_1 z = z = T_2 z$ . Thus  $z$  is the common fixed point of pair of maps  $T_1$  and  $T_2$ . This completes the proof.

**Theorem 3.5:** Let  $(X, k)$  be a digital image where  $X \subseteq Z^n$  and  $k$  is an adjacency relation in  $X$ . Let  $(X, d, k)$  be a complete digital metric space and suppose  $T_1, T_2 : X \rightarrow X$  be any two continuous and onto mapping satisfying the condition

$$\begin{aligned}
 d(T_1 x, T_2 y) & \geq \alpha \max\{d(x, y), d(x, T_1 x), d(y, T_2 y)\} \\
 & + \beta \max\{d(x, T_2 y), d(x, y)\} + \gamma d(x, y)
 \end{aligned}$$

for all  $x, y \in X$ , where  $\alpha \geq 0, \beta > 0, \gamma \leq 1$  are constant, with  $\alpha + \beta + \gamma > 1$ . Then  $T_1$  and  $T_2$  have a common fixed point in  $X$ .

**Proof:** Let  $x_0$  be an arbitrary point in  $X$ . Since  $T_1$  and  $T_2$  be onto (surjective), there exist  $x_0 \in X$  and  $x_1 \in X$



such that

$$T_1(x_1) = x_0, T_2(x_2) = x_1$$

In this way, we define the sequences  $\{x_{2n}\}$  and  $\{x_{2n+1}\}$  by

$$x_{2n} = T_1x_{2n+1} \text{ for } n = 0,1,2,3 \dots \dots \text{and}$$

$$x_{2n+1} = T_2x_{2n+2} \text{ for } n = 0,1,2,3 \dots \dots$$

Note that, If  $x_{2n} = x_{2n+1}$  some  $n \geq 1$ , then it is fixed point of  $T_1$  and  $T_2$ .

Now putting  $x = x_{2n+1}$  and  $y = x_{2n+2}$ , we have

$$d(x_{2n}, x_{2n+1}) = d(T_1x_{2n+1}, T_2x_{2n+2})$$

$$\begin{aligned} \Rightarrow d(x_{2n}, x_{2n+1}) &\geq \alpha \max \{d(x_{2n+1}, x_{2n+2}), d(x_{2n+1}, T_1x_{2n+1}), d(x_{2n+2}, T_2x_{2n+2})\} \\ &\quad + \beta \max \{d(x_{2n+1}, T_2x_{2n+2}), d(x_{2n+1}, x_{2n+2})\} + \gamma d(x_{2n+1}, x_{2n+2}) \\ &\geq \alpha \max \{d(x_{2n+1}, x_{2n+2}), d(x_{2n+1}, x_{2n}), d(x_{2n+2}, x_{2n+1})\} \\ &\quad + \beta \max \{d(x_{2n+1}, x_{2n+1}), d(x_{2n+1}, x_{2n+2})\} + \gamma d(x_{2n+1}, x_{2n+2}) \end{aligned}$$

$$\Rightarrow d(x_{2n}, x_{2n+1}) \geq (\alpha + \beta + \gamma) d(x_{2n+1}, x_{2n+2})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq \frac{1}{(\alpha + \beta + \gamma)} d(x_{2n}, x_{2n+1})$$

$$\Rightarrow d(x_{2n+1}, x_{2n+2}) \leq h d(x_{2n}, x_{2n+1}), \text{ where } h = \frac{1}{(\alpha + \beta + \gamma)}, 0 \leq h \leq 1.$$

In general

$$d(x_{2n}, x_{2n+1}) \leq h d(x_{2n-1}, x_{2n}) \leq \dots, \dots, \dots h^{2n} d(x_0, x_1)$$

So for  $n < m$ , we have

$$\begin{aligned} d(x_{2n}, x_{2m}) &\leq d(x_{2n}, x_{2n+2}) + \dots, \dots, \dots + d(x_{2m-1}, x_{2m}) \\ &\leq (h^{2n} + h^{2n+1} + \dots + h^{2m-1}) d(x_0, x_1) \\ &\leq \frac{h^{2n}}{1-h} d(x_0, x_1) \end{aligned}$$

Let  $0 \leq c$  be given, choose a natural number  $N_1$  such that  $\frac{h^{2n}}{1-h} d(x_0, x_1) \leq c$ , for all  $n \geq N_1$ . Thus  $d(x_{2n}, x_{2m}) \leq c$ , for  $n < m$ . Therefore  $\{x_{2n}\}$  is a Cauchy sequence in  $(X, d, k)$ . Since  $(X, d, k)$  is a complete digital metric space, there exists  $z \in X$  such that  $x_{2n} \rightarrow z$  as  $n \rightarrow \infty$ . If  $T_1$  is continuous, then

$$d(T_1 z, z) \leq d(T_1 x_{2n+1}, T_1 z) + d(T_1 x_{2n+1}, z) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Since  $x_{2n} \rightarrow z$  and  $T_1 x_{2n+1} \rightarrow T_1 z$  as  $n \rightarrow \infty$ . Therefore  $d(T_1 z, z) = 0$ . This implies that,  $T_1 z = z$ . Hence  $z$  is a fixed point of  $T_1$ .

Similarly, it can be established that  $T_2 z = z$ . Therefore,  $T_1 z = z = T_2 z$ . Thus  $z$  is the common fixed point of pair of maps  $T_1$  and  $T_2$ . This completes the proof.

**CONCLUSION**

Since a digital images be a combination of pixels in 2-D while voxels are in 3-D. And distance function between these pixels may be contractive or expansive. On this concept to find the diversified

image we took two self-mapping over complete digital metric space via five various types of expansive conditions and proved fixed point theorems. We hope these theorems provide a new dimension of very traditional FPT to the modern theory of Image processing.

## REFERENCES

- [1]. Dolhare, U.P. and Nalawade, V.V. (2018), "Fixed point theorems in digital images and applications to fractal image compression." Asian Journal of Mathematics and Computer Research; Vol. 25(1), pp. 18-37.
- [2]. Edelstein, M. (1961), "An extension of Banach contraction principle." Proceeding American Mathematical Society; Vol. 12(1), pp. 7-10.
- [3]. Gupta, N., Singh, A. and Modi, G. (2019), "Application of fixed point theorems for digital contractive type mappings in digital metric spaces." International Journal of Scientific Research in Mathematical and Statistical Sciences, E-ISSN: 2348-4519; Vol. 6, Issue-1, pp. 323-327.
- [4]. Han, S.E. (2016), "Banach fixed point theorem from the viewpoint of digital topology." Journal non linear Science Applications, Vol. 9, pp. 895-905.
- [5]. Jain, D. (2018), "Common fixed point theorem for intimate mappings in digital metric spaces." International Journal of Mathematics Trends and Technology; Vol. 56 (2), pp. 91-94.
- [6]. Jyoti, K. and Rani, A. (2017), "Digital expansions endowed with fixed point theory." Turkish Journal of Analysis and Number Theory, Vol. 5(5), PP 146-152.
- [7]. Mishra, A., Tripathi, P.K., Agrawal, A.K. and Joshi, D.R. (2019), "A contraction mapping method in digital image processing." International Journal of Recent Technology and Engineering, ISSN: 2277-3878, Vol. 8, Issue-455
- [8]. Mishra, L.N., Jyoti, K., Rani, A. and Vandana (2018), "Fixed point Theorems with digital Contractions image processing." Non linear Science letter, Vol. 9(2), pp. 104-115.
- [9]. Rani, A. and Jyoti, K. (2016), "Common fixed point theorems in digital metric spaces." International Journal of Scientific & Engineering Research, ISSN: 2229-5518, Vol. 7, Issue 12, PP 1704-1716.
- [10]. Rosenfeld, A. (1979), "Digital Topology." American Mathematical Society; Vol. 86, pp. 621-630.
- [11]. Sridevi, K., Kiran, D.M.K., and Kameswari, M.V.R. (2017), "Fixed point theorems for digital contractive type mappings in digital metric spaces." International Journal of Mathematics Trends and Technology; Vol. 48 (3), pp. 159-167.
- [12]. Suganthi, A. (2021), "Common fixed point theorems in digital metric space using implicit relation." Advances and Application in Mathematical Sciences; Vol. 20 Issue 10; PP 2351-2369.