



<http://www.bomsr.com>

Email:editorbomsr@gmail.com

RESEARCH ARTICLE



A Modish Glimpse on Mag-Shanthi Integer Sequence

J. Shanthi¹, M.A. Gopalan²

¹ Assistant Professor, Department of Mathematics, Shrimati Indira Gandhi College, Affiliated to Bharathidasan University, Trichy-620 002, Tamil Nadu, India.

Email: shanthivishvaa@gmail.com,

² Professor, Department of Mathematics, Shrimati Indira Gandhi College, Affiliated to Bharathidasan University, Trichy-620 002, Tamil Nadu, India.

Email: mayilgopalan@gmail.com

DOI:10.33329/bomsr.11.2.38



J. Shanthi

ABSTRACT

A new integer sequence called Mag-Shanthi integer sequence generated from the recurrence relation $MS_{n+2} = 8MS_{n+1} - 7MS_n, n \geq 0$ with the initial conditions $MS_0 = 6, MS_1 = 48$ is analyzed for varieties of interesting properties.

Keywords: integer sequence, Mag-Shanthi numbers

Introduction:

Number is the essence of mathematical calculations. Numbers have varieties of patterns and have varieties of range and richness. Many numbers exhibit fascinating properties, they form sequences, they form patterns and so on. In this context one may refer [1-12]. This communication presents varieties of fascinating properties on Mag-Shanthi numbers.

Method of analysis :

The Mag-Shanthi sequence denoted by $\{MS_n\}$ is defined by the recurrence relation

$$MS_{n+2} = 8MS_{n+1} - 7MS_n \quad (1)$$

with the initial conditions

$$MS_0 = 6, MS_1 = 48 \quad (2)$$

The auxiliary equation associated with the recurrence relation (1) is given by

$$m^2 - 8m + 7 = 0 \quad (3)$$

whose roots are

$$m_1 = 1, m_2 = 7$$

Thus, the general solution of (1) is

$$MS_n = A7^n + B$$

From the initial conditions, we infer that

$$A = 7, B = -1$$

Hence, the n^{th} integer in the sequence $\{MS_n\}$ is given by

$$MS_n = 7^{n+1} - 1$$

A few interesting properties among Mag-Shanthi numbers are presented below :

Property 1: $MS_{n+1} - 7MS_n = 6$

Property 2: $288(MS_{n+1} - 7MS_n) + 1 = 1729$, Taxicab number

Property 3: $6(MS_{2n+1} - 2MS_{2n} + MS_{2n-1}) = \text{area of pythagorean triangle } (18 * 7^n, 24 * 7^n, 30 * 7^n)$

Proof:

$$\begin{aligned} \text{L.H.S.} &= 6[(7^{2n+2} - 1) - 2(7^{2n+1} - 1) + (7^{2n} - 1)] \\ &= 6(6 * 7^n)^2 \\ &= \text{area of Pythagorean triangle } (18 * 7^n, 24 * 7^n, 30 * 7^n) \end{aligned}$$

Property 4: $MS_{k+n+k-1} = 7MS_{k+n+k-2} + 6$

Proof:

$$\begin{aligned} \text{L.H.S.} &= 7^{k+n+k} - 1 = 7(7^{k+n+k-1} - 1 + 1) - 1 \\ &= 7(7^{k+n+k-1} - 1) + 6 = \text{R.H.S.} \end{aligned}$$

Property 5: $MS_{3n+2} = MS_n * MS_{2n+1} + MS_n + MS_{2n+1}$

Proof :

$$\begin{aligned} \text{L.H.S.} &= 7^{3n+3} - 1 = 7^{n+1} * 7^{2n+2} - 1 \\ &= (7^{n+1} - 1)(7^{2n+2} - 1) + 7^{n+1} + 7^{2n+2} - 2 \\ &= (7^{n+1} - 1)(7^{2n+2} - 1) + (7^{n+1} - 1) + (7^{2n+2} - 1) \\ &= \text{R.H.S.} \end{aligned}$$

Property 6: $\sum_{i=1}^n MS_i = \frac{7MS_n - 6(7+n)}{6}$

Proof:

$$\begin{aligned} \text{L.H.S.} &= \sum_{i=1}^n (7^{i+1} - 1) = 7 \sum_{i=1}^n 7^i - n = \frac{7(7^{n+1} - 7)}{6} - n \\ &= \frac{7(7^{n+1} - 1 - 6)}{6} - n = \frac{7MS_n}{6} - 7 - n = \text{R.H.S.} \end{aligned}$$

Property 7: $\sum_{i=1}^n (MS_i)^2 = \frac{49MS_{2n+1} - 112MS_n + 48(n-35)}{48}$

Proof:

$$\begin{aligned} \text{L.H.S.} &= 7^2 \sum_{i=1}^n 7^{2i} - 14 \sum_{i=1}^n 7^i + n \\ &= \frac{7^2(7^{2n+2} - 49)}{48} - \frac{14(7^{n+1} - 7)}{6} + n \\ &= \frac{7^2(MS_{2n+1} - 48)}{48} - \frac{7(MS_n - 6)}{3} + n \\ &= \frac{49MS_{2n+1}}{48} - \frac{7MS_n}{3} - 35 + n \\ &= \text{R.H.S.} \end{aligned}$$

Property 8: $MS_p * MS_q = 7MS_{p+q} - MS_p - MS_q + 6$

Proof:

$$\begin{aligned} \text{L.H.S.} &= (7^{p+1} - 1) * (7^{q+1} - 1) = 7^{p+q+2} - 7^{p+1} - 7^{q+1} + 1 \\ &= 7(7^{p+q+1} - 1 + 1) - (7^{p+1} - 1 + 1) - (7^{q+1} - 1 + 1) + 1 \\ &= 7MS_{p+q} - MS_p - MS_q + 6 = \text{R.H.S.} \end{aligned}$$

Property 9: $\sum_{i=0}^n (MS_{i+1} - MS_i) = 7MS_n$

Proof:

$$\begin{aligned} \text{L.H.S.} &= (MS_1 - MS_0) + (MS_2 - MS_1) + \dots + (MS_n - MS_{n-1}) + (MS_{n+1} - MS_n) \\ &= MS_{n+1} - MS_0 = (7^{n+2} - 1) - 6 \\ &= 7^{n+2} - 7 = 7(7^{n+1} - 1) = 7 MS_n = \text{R.H.S.} \end{aligned}$$

Property 10: $MS_n \left(\sum_{i=1}^k 7^i (MS_{(i+1)n} + 1) \right) = (MS_n + 1)^2 ((MS_n + 1)^k - 1)$

Proof:

By definition ,one has

$$MS_n + 1 = 7^{n+1} = 7 * 7^n \tag{4}$$

Replacing n by 2 n in (4) , we have

$$MS_{2n} + 1 = 7 * 7^{2n} \tag{5}$$

From (4) & (5) ,one obtains after some algebra that

$$7 (MS_{2n} + 1) = (MS_n + 1)^2 \tag{6}$$

Replacing n by 3 n ,4 n ,5 n ,....in (4) , it is seen in general that

$$7^i (MS_{(i+1)n} + 1) = (MS_n + 1)^{i+1}, i = 1,2,3,...$$

Therefore ,

$$\sum_{i=1}^k 7^i (MS_{(i+1)n} + 1) = \sum_{i=1}^k (MS_n + 1)^{i+1} \tag{7}$$

Observe that

$$\sum_{i=1}^k (MS_n + 1)^{i+1} = \frac{(MS_n + 1)^2 ((MS_n + 1)^k - 1)}{MS_n} \tag{8}$$

Substituting (8) in (7) ,the required result is obtained.

Property 11: $48 \sum_{i=1}^k (MS_i + 1)^2 = 7^4 MS_{2k-1}$

Proof:

Consider (6) of Property 10. We have

$$\begin{aligned}
\sum_{n=1}^k (MS_n + 1)^2 &= 7 \sum_{n=1}^k (MS_{2n} + 1) \\
&= 7 \sum_{n=1}^k MS_{2n} + 7k \\
&= 7 \sum_{n=1}^k (7^{2n+1} - 1) + 7k \\
&= 7^2 \sum_{n=1}^k 7^{2n} \\
&= 7^4 \frac{(7^{2k} - 1)}{(7^2 - 1)} \\
&= 7^4 \frac{MS_{2k-1}}{48}
\end{aligned}$$

Hence the property.

Property 12: $MS_{k_{n+1}} = 7^2 MS_{k_{n-1}} + 48$

Proof:

$$\begin{aligned}
MS_{k_{n+1}} &= 7^{k_{n+2}} - 1 = 7^2 (7^{k_{n-1+1}} - 1 + 1) - 1 \\
&= 7^2 MS_{k_{n-1}} + 7^2 - 1 = 7^2 MS_{k_{n-1}} + 48
\end{aligned}$$

Property 13: $7(MS_{2n+1} + 1) - (MS_n + 1)^2$ is a square multiple of 6

Proof:

$$7(MS_{2n+1} + 1) - (MS_n + 1)^2 = 7 * 7^{2n+2} - (7^{n+1})^2 = 6 * (7^{n+1})^2$$

Property 14: $\sum_{i=1}^n MS_i * MS_{i+1} = \frac{7^3 MS_{2n+1}}{48} - \frac{28 MS_n}{3} + n - 287$

Proof:

$$\begin{aligned}
\text{L.H.S.} &= \sum_{i=1}^n (7^{i+1} - 1)(7^{i+2} - 1) \\
&= 7^3 \sum_{i=1}^n 7^{2i} - 56 \sum_{i=1}^n 7^i + n \\
&= 7^3 \left[\frac{7^2(7^{2n} - 1)}{48} \right] - 56 \left[\frac{7(7^n - 1)}{6} \right] + n \\
&= \frac{7^3}{48} (MS_{2n+1} - 48) - \frac{28}{3} (MS_n - 6) + n \\
&= \frac{7^3}{48} MS_{2n+1} - \frac{28}{3} MS_n + n - 287 = \text{R.H.S.}
\end{aligned}$$

For simplicity and brevity ,some more properties satisfied by Mag-Shanthi numbers are exhibited below:

$$\text{Property 15: } \sum_{i=0}^{n-1} (MS_i + 1)(MS_i + 2) = \frac{49*7^{2n} + 56*7^n - 105}{48}$$

Property 16 : $(MS_n + 1) (MS_n + 2)$ is twice the Triangular number of rank 7^{n+1}

Property 17 : $(MS_n + 1) (MS_n + 2) (MS_n + 3)$ is six times the Triangular pyramidal number of rank 7^{n+1}

Property 18 : $(MS_n + 1)^2 (MS_n + 2)$ is twice the Pentagonal pyramidal number of rank 7^{n+1}

Property 19 : Powers of Mag-Shanthi numbers are presented :

$$(MS_n)^2 = MS_{2n+1} - 2MS_n$$

$$(MS_n)^3 = MS_{3n+2} - 3MS_{2n+1} + 3MS_n$$

$$(MS_n)^4 = MS_{4n+3} - 4MS_{3n+2} + 6MS_{2n+1} - 4MS_n$$

$$(MS_n)^5 = MS_{5n+4} - 5MS_{4n+3} + 10MS_{3n+2} - 10MS_{2n+1} + 5MS_n$$

$$(MS_n)^6 = MS_{6n+5} - 6MS_{5n+4} + 15MS_{4n+3} - 20MS_{3n+2} + 15MS_{2n+1} - 6MS_n$$

and so on. Observe the beautiful pattern of coefficients in the above representations.

Property 20 : $14*$ Pentagonal pyramidal number of rank $7^n - MS_{n-1} + MS_{2n-1} (1 - MS_n) + 2$ is a perfect square.

Conclusion

In this paper , a new sequence of integers named as Mag-Shanthi numbers have been introduced along with interesting relations among Mag-Shanthi numbers. One may search for other connections between Mag-Shanthi numbers.

References

- [1]. M.A.Gopalan, A. Gnanam, "A Notable Integer Sequence ", Mathematical Sciences Research Journal an International journal Rapidpublication, Volume 12, Number 1, Pp 7-15, January 2008.
- [2]. M.A. Gopalan, G. Srividhya, " Higher ordercongruent numbers", Pacific – Asian Journal of Mathematics, Volume 2, No. 1-2, Pp 63-78, January-December 2008.
- [3]. Dr.M.A.Gopalan, A. Gnanam, "Star numbers", Mathematical Sciences Research Journal, 12(12), Pp 303-308, 2008.
- [4]. M.A.Gopalan, N. Vanitha and Manju Somanath, " Observations on Carmichael numbers with four factors", Bulletin of Pure and Applied Sciences, Volume 27, E(No.1), Pp 17-19, 2008.
- [5]. M.A.Gopalan, G. Srividhya Krishnamoorthy, " On congruent numbers", Bulletin of Pure and applied Sciences, Volume 28, E(No.1), Pp 31-40, 2009.
- [6]. M.A.Gopalan, A. Gnanam, " Shrodder numbers", Proceedingof the International Conference on Mathematical Methods and Computation, 24-25, Pp 84-86, July 2009.
- [7]. M.A.Gopalan, A. Gnanam, " Kynea numbers", Bulletin of Pure and Applied Sciences, Vol. 29, E(No.1), Pp 73-76, 2010.

- [8]. M.A. Gopalan , A.Gnanam, “ Four Dimensional Pyramidal numbers”, Pacific-Asian Journal of Mathematics, Volume 4, No. 1, Pp 53-62, January-June 2010.
- [9]. M.A. Gopalan, K. Geetha, Manju Somanath, “ Observations on Icosahedral numbers”, International journal of Engineering Research –Online , A peer Reviewed international journal, Vol. 1., Issue.3., Pp 395-400.
- [10]. M. A. Gopalan, S. Vidhyalakshmi, J. Shanthi, “ A new Integer Sequence”, International Journal of Recent Trends in Engineering and Research, Volume-2, Issue-6, June-2016, 307-314.
- [11]. M. A. Gopalan, S. Vidhyalakshmi, J. Shanthi, “ A Remarkable Integer Sequence”, International Journal of Engineering & Scientific Research, Volume-4, Issue-8, August-2016, 37-45.
- [12]. J.Shanthi,S.Vidhyalakshmi,M.A.Gopalan,New Vistas on Mersenne numbers, The Ciencia & Engenharia-Science &Engineering Journal,Volume-11,Issue-1,2023,356-363