



http://www.bomsr.com  
Email:editorbomsr@gmail.com

**RESEARCH ARTICLE**

*A Peer Reviewed International Research Journal*



**CLASSIFICATIONS IN FUZZY PSEODOMETRIC SPACES**

**SERVET KUTUKCU<sup>1\*</sup>, ELIF AYDIN<sup>2</sup>**

<sup>1,2</sup>Department of Mathematics, OndokuzMayis University 55139 Kurupelit, Samsun, Turkey

\*E-mail: skutukcu@omu.edu.tr



**ABSTRACT**

In this paper, we give strong and stationary structures in fuzzy pseodometric spaces and present relations between them illustrating examples.

Keywords: strong, stationary, fuzzy pseodometric.

AMS(2010) Subject Classification: 54A40, 54E35.

**©KY PUBLICATIONS**

**1. INTRODUCTION**

A triangular norm, shortly t-norm, is a kind binary operation on the unit interval [0,1] used in multivalued logic especially fuzzy logic generalizes intersection in a lattice, conjunction in logic and triangle inequality in ordinary metric spaces. In [6], a t-norm  $*$  : [0,1] × [0,1] → [0,1] defined as  $a*b=$  for all  $a∈[0,1]$ ,  $*$  is symmetric,  $*$  is nondecreasing in each variable and  $*$  is associative. We will make use of three basic t-norms, namely the minimum operator, the algebraic product and the Lukasiewicz t-norm TL defined by  $a*b = \min\{a,b\} = TM$ ,  $a*b = a.b = TP$  and  $a*b = \max\{0, a+b-1\} = TL$ , respectively. These t-norms are ranked as  $TL \leq TP \leq TM$ , in fact, TM is the strongest t-norm. Using t-norms to generalize triangle inequality, George and Veeramani introduced fuzzy metric space,

**Definition 1.1 [1].** A 3-tuple  $(X,M,*)$  is called a fuzzy metric space, shortly FM-space, if X is an arbitrary set,  $*$  is a continuous t-norm and M is a fuzzy set on  $X^2 \times (0,\infty)$  satisfying following conditions; for all  $x,y,z \in X$  and  $t,s > 0$

- FM1)  $M(x,y,t) > 0,$
- FM2)  $M(x,y,t) = 1$  iff  $x = y,$
- FM3)  $M(x,y,t) = M(y,x,t),$
- FM4)  $M(x,y,t+s) \geq M(x,z,t) * M(z,y,s),$
- FM5)  $M(x,y,.) : (0,\infty) \rightarrow [0,1]$  is continuous.

Also M is called a fuzzy metric on X.

If we take  $FM2^*$   $x = y$  then  $M(x,y,t) = 1$  instead of FM2, then  $(X,M,*)$  is called fuzzy pseudometric space, shortly FPM-space.

If we take minimum operator  $*$ , then  $(X,M,*)$  is called fuzzy ultrametric space.

Throughout the paper, we will denote  $(0,\infty)$  with  $\mathbb{R}^+$ .

## 2.STRONG FUZZY PSEODOMETRIC SPACES

**Definition 2.1.** Let  $(X,M,*)$  be a FPM-space.  $M$  is said to be a strong fuzzy pseudometric on  $X$ , or  $(X,M,*)$  is said to be a strong FPM-space if  $M(x,y,t) \geq M(x,z,t) * M(z,y,t)$  for all  $x,y,z \in X$  and  $t > 0$ .

**Remark 2.1.** From Definition 1.1, every strong FPM-space is a FPM-space, but the converse is not true.

**Example 2.1.** Let  $X = \{x,y,z\}$  and  $a*b = a.b$  for all  $a,b \in [0,1]$ . Define  $M: X \times X \times (0,\infty) \rightarrow [0,1]$  by  $M(x,x,t) = M(y,y,t) = M(z,z,t) = 1$ ,  $M(x,z,t) = M(z,x,t) = M(y,z,t) = M(z,y,t) = t/t+1$  and  $M(x,y,t) = M(y,x,t) = t^2/(t+2)^2$ . Then  $M$  is a fuzzy pseudometric metric but not a strong fuzzy pseudometric on  $X$ .

**Example 2.2.** Let  $X = \{x,y,z\}$  and  $a*b = \max\{0,a+b-1\}$  for all  $a,b \in [0,1]$ . Define  $M: X \times X \times (0,\infty) \rightarrow [0,1]$  by  $M(x,x,t) = M(y,y,t) = M(z,z,t) = 1$ ,  $M(x,z,t) = M(z,x,t) = M(y,z,t) = M(z,y,t) = 2t+1/2t+2$  and  $M(x,y,t) = M(y,x,t) = t/t+2$ . Then  $M$  is a fuzzy pseudometric metric but not a strong fuzzy pseudometric on  $X$ .

**Example 2.3.** Let  $X = \mathbb{R}^+$  and  $a*b = a.b$  for all  $a,b \in [0,1]$ . Define  $M: X \times X \times (0,\infty) \rightarrow [0,1]$  by  $M(x,y,t) = \min\{x,y\}/\max\{x,y\}$ . Then  $M$  is a fuzzy pseudometric metric and also a strong fuzzy pseudometric on  $\mathbb{R}^+$ .

**Theorem 2.1.** Let  $(X,d)$  be a pseudometric space and  $a*b = a.b$  for all  $a,b \in [0,1]$ . Define  $M_d: X \times X \times (0,\infty) \rightarrow [0,1]$  by  $M_d(x,y,t) = t/t+d(x,y)$ . Then  $M_d$  is a fuzzy pseudometric metric and also a strong fuzzy pseudometric on  $X$ .  $(X,M_d,*)$  is called standard strong FPM-space.

**Proof.** In [8], we know that  $(X,M_d,*)$  is a FPM-space. We only show that  $M_d$  is strong. Since  $(X,d)$  is a pseudometric space, we have  $d(x,z) \leq d(x,y) + d(y,z)$  for  $x,y,z \in X$ . Then

$$\begin{aligned} 1 + \frac{d(x,z)}{t} &\leq 1 + \frac{d(x,y) + d(y,z)}{t} \\ &\leq \frac{t^2 + td(x,y) + td(y,z) + d(x,y).d(y,z)}{t^2} \\ &\leq \frac{[t + d(x,y)][t + d(y,z)]}{t^2} \end{aligned}$$

and so,

$$\frac{t}{t + d(x,z)} \geq \frac{t}{t + d(x,y)} \cdot \frac{t}{t + d(y,z)}$$

hence,

$$M_d(x,z,t) \geq M(x,y,t) * M(y,z,t).$$

This completes the proof.

**Definition 2.2.** Let  $(X,M,*)$  be a strong FPM-space. If  $*$  is minimum operator then  $(X,M,*)$  is said to be a strong fuzzy ultra-pseudometric space.

The proof of following theorem is easily omitted.

**Theorem 2.2.** Standard strong FPM-space  $(X,M_d,*)$  is a strong fuzzy ultra-pseudometric space if and only if  $d$  is ultra-pseudometric on  $X$ .

### 3. STATIONARY FUZZY PSEODOMETRIC SPACES

**Definition 3.1.** Let  $(X, M, *)$  be a FPM-space.  $M$  is said to be a stationary fuzzy pseudometric on  $X$  or  $(X, M, *)$  is said to be a stationary FPM-space if  $M$  does not depend on  $t$ , i.e.  $M(x, y, t) = M(x, y)$  is constant for all  $x, y \in X$ .

**Remark 3.1.** From Definition 1.1, every stationary FPM-space is a FPM-space, but the converse is not true.

**Example 3.1.** From [8], we know that  $(X, M_d, *)$  is a FPM-space. But  $(X, M_d, *)$  is not a stationary FPM-space since the function  $M_d$  depends on  $t$ . If we take  $t = 1$ , then  $(X, M_d, *)$  is a stationary FPM-space. For  $t = 1$ , we call  $(X, M_d, *)$  is a standard stationary FPM-space.

**Example 3.2.** A FPM-space  $(X, M, *)$  give in Example 2.3 is a stationary FPM-space.

**Example 3.3.** Let  $X = (0, 1/2)$  and  $a * b = \max\{0, a + b - 1\}$  for all  $a, b \in [0, 1]$ . Define  $M: X \times X \times (0, \infty) \rightarrow [0, 1]$  by

$$M(x, y, t) = \begin{cases} 1, & x = y \\ x + y, & x \neq y \end{cases}. \text{ Then } M \text{ is a stationary fuzzy pseudometric on } X.$$

**Theorem 3.1.** Every stationary FPM-space is a strong FPM-space.

**Proof.** Let  $(X, M, *)$  be a stationary FPM-space. Then, from (FM4),  $M(x, y) \geq M(x, z) * M(z, y)$  for all  $x, y \in X$ . Since  $M$  is not depend on  $t$ , we have  $M(x, y, t) \geq M(x, z, t) * M(z, y, t)$ . From Definition 2.1,  $(X, M, *)$  is a strong FPM-space.

This completes the proof.

**Remark 3.2.** The converse of Theorem 3.1 is not true.

**Example 3.4.** Let  $X = \mathbb{R}^+$  and  $a * b = \min\{a, b\}$  for all  $a, b \in [0, 1]$ . Define  $M: \mathbb{R}^+ \times \mathbb{R}^+ \times (0, \infty) \rightarrow [0, 1]$  by

$$M(x, y, t) = \begin{cases} 1, & x = y \\ \frac{t}{t+1}, & x \neq y \end{cases}. \text{ Then } M \text{ is a strong fuzzy pseudometric but not a stationary fuzzy}$$

pseudometric on  $X$ .

**Theorem 3.2.** Let  $(X, M, *)$  be a FPM-space and define a function family  $\{M_t : t \in \mathbb{R}^+\}$  as  $M_t: X \times X \rightarrow (0, 1]$  by  $M_t(x, y) = M(x, y, t)$ . Then  $(X, M_t, *)$  is a stationary FPM-space for all  $t \in \mathbb{R}^+$  if and only if  $(X, M, *)$  is a strong FPM-space.

**Proof.** Let  $(X, M_t, *)$  be a stationary FPM-space for all  $t \in \mathbb{R}^+$ . Then, from Theorem 3.1,  $(X, M_t, *)$  is a strong FPM-space for all  $t \in \mathbb{R}^+$ . Since  $M_t(x, y) = M(x, y, t)$  for all  $t \in \mathbb{R}^+$ , then  $(X, M, *)$  is a strong FPM-space.

Let  $(X, M, *)$  be a strong FPM-space. Then, from Remark 3.1,  $(X, M, *)$  is a FPM-space. Define a function  $\varphi: \mathbb{R}^+ \rightarrow (0, 1]$  by  $\varphi(t) = k$  (a constant), then  $\varphi$  is increasing, continuous and

$M(x, y, t) = M(x, y, k) = M_k(x, y) = M_t(x, y)$ . So the distance  $M(x, y, t)$  is not depend on  $t$  for all  $t > 0$ . Hence,  $(X, M, *)$  i.e.  $(X, M_t, *)$  be a stationary FPM-space for all  $t \in \mathbb{R}^+$ .

This completes the proof.

**Definition 3.2.** Let  $(X, M, *)$  be a stationary FPM-space. If  $*$  is minimum operator then  $(X, M, *)$  is said to be a stationary fuzzy ultra-pseudometric space.

The proof of following theorem is easily omitted.

**Theorem 3.3.** Standard stationary FPM-space  $(X, M_d, *)$  is a stationary fuzzy ultra-pseudometric space if and only if  $d$  is ultra-pseudometric on  $X$ .

### 3. OPEN PROBLEMS

**Problem 3.1.** In [3], Kutukcu et.al defined intuitionistic fuzzification of Menger spaces. What are intuitionistic classifications of the notions strong and stationary?

**Problem 3.2.** In [4], Park defined intuitionistic fuzzification of metric spaces. What are intuitionistic classifications of the notions strong and stationary?

#### 4. REFERENCES

- [1]. George A., Veeramani P., 1994, On some results in fuzzy metric spaces, *Fuzzy Sets Syst.* 64, 395-399.
  - [2]. Gregori, V., Morillas, S., Sapena, A., 2010, On A Class of Completable Fuzzy Metric Spaces, *Fuzzy Sets Syst.* 161, 2193 – 2205.
  - [3]. Kutukcu S., Tuna A., Yakut A.T., 2007, Generalized contraction mapping principle in intuitionistic Menger spaces and application to differential equations, *Appl. Math. Mech. (English Ed.)* 12, 799-809.
  - [4]. Park J.H., 2004, Intuitionistic fuzzy metric spaces, *Chaos, Solitons Fractals* 22, 1039-1046.
  - [5]. Sapena, A., Morillas, S., WIAT'09, On strong fuzzy metrics, *Proc. Workshop in Applied Topology*, 135 – 141.
  - [6]. Schweizer B., Sklar A., 1960, Statistical metric spaces, *Pacific Journal Math.*, 10, 314-334.
-