



<http://www.bomsr.com>

Email:editorbomsr@gmail.com

RESEARCH ARTICLE

A Peer Reviewed International Research Journal




DYNAMICAL SYSTEMS ARISING FROM VECTOR FIELDS ON SMOOTH MANIFOLDS

Dr (Smt). S. N. BANASODE¹, SEEMA B PATIL², JAYASHREE .S. RADDER³

¹Associate Professor of Mathematics, R.L Science Institute Belagavi, Karnataka State, India.

²Research Scholar, Rani Channamma University Belagavi. Karnataka State, India.

³Assistant Professor of Mathematics, Tontadarya College of Engg,
Gadag, Karnataka State, India.



ABSTRACT

In this paper we have considered a problem that looks into a possible lift for functions defined on manifolds and flows associated with the dynamical systems on them. Initially we observe this problem on one dimensional manifolds. Later ,it was generalized to M where n is a dimensional smooth manifold $\mathbb{R}^n, n > 1$.

Keywords: Smooth manifolds ,vector fields, orientable manifolds.

©KY PUBLICATIONS

1. INTRODUCTION

In this section we give account of smooth differentiable manifolds which are finite dimensional. Non trivial examples of manifolds which admits smooth differentiable structures are S^1 the circle group,also denoted by T^1 the one dimensional torus.They generally come from quotient topology induced by topology of R^2 .The Cartesian product of the circle group taken n times gives rise to n torus which is also regarded as quotient space with induced topology of R^{n+1} .Our manifolds are orientable manifolds and the maps are orientation preserving maps ,quite naturally such Orientation preserving homeomorphisms give rise to dynamical systems and if the maps are smooth this dynamical systems are also smooth.. The theory of dynamical systems is in fact vast area of study initiated by Henri Poincare around 1900. He was interested studying problems in celestial mechanics. Later Kolomogrov ,Birkoff ,Arnold, Kac and many others have continued to investigate the problems arising in this area of research.In this paper We have noticed that a vector field arising from an integral curve giving rise to. a dynamical system which is a system of first order ordinary differential equations .

1.1 Vector fields as Flows

Let M be a smooth n -dimensional manifolds then it admits a smooth differentiable structure.

Let $x \in M$ lying in some coordinate patch of (U_i) of M where U is an open subset of M and i , $U:U \rightarrow \mathbb{R}^n$ is a local diffeomorphism.

Let $T_x M$ denote the tangent space of M at x then we know that $T_x M \sim \mathbb{R}^n$

i.e there is an isomorphism with local basis with vector fields

$\{\partial/\partial x_i : i = 1, 2, \dots, n\}$ of $T_x M$. To this end consider a smooth curve $\gamma: I \rightarrow M$,

I is the interval containing zero such that $\gamma(0) = x$

$t \rightarrow \gamma(t) \quad t \in I$

then $\gamma^1(t) = \frac{d\gamma(t)}{dt}$ is the velocity vector

In fact a tangent vector to the curve at $\gamma(t)$ in M .

Since $\gamma^1(t)$ is a tangent vector to M at $\gamma(t)$ i.e $\gamma^1(t) \in T_{\gamma(t)} M$

We have proved the following proposition .

Proposition 1: If $\gamma: I \rightarrow M$ is a smooth curve passing through x in M , I containing zero then $\gamma^1(t) \in T_{\gamma(t)} M$, where $T_{\gamma(t)} M$ is the tangent space of M at $\gamma(t)$.

In particular $\gamma^1(0)$ is the tangent vector to M at x of $T_{\gamma(0)} M$. This been a local description vector field on M provided a global picture because vector field in M is a map.

$X: M \rightarrow TM$ where TM is the tangent bundle of M (this bundle is smooth because M is smooth) which attaches to each point P of M then the tangent vector $X(P)$.

$P \rightarrow X(P)$

Here $TM = \cup_{\gamma(t)} T_{\gamma(t)} M$

In other words this is more so of defining a vector in an invariant fashion as a map

$X: M \rightarrow TM$

Such that $\pi \circ X = \text{Id}_M$ where $\pi: TM \rightarrow M$ is a projection maps.

Observation:

The map that assigns to each point of M at tangent vector gives rise to a system of first order ordinary differential equation.

$X(P): \frac{dX(P)}{dP}$

1.2 Dynamical Systems:

We shall set up a simple dynamical system

$X^1(t) = rX(t)$, where X is a smooth vector field in M , where X is a smooth vector field in M .

where $r > 0$ is some constant.

$f: \mathbb{R} \rightarrow \mathbb{R} \quad x^1 = ax, \quad x(0) = a$

$x(t) = ae^{at}$

$x^1(t) = ae^{at}$

i.e $= ax(t)$

$x(0) = e^{a0} = 1$

$x^1(t) = x(0)x(t)$

$\frac{x^1(t)}{x(t)} = x(0)$

Some digression

Let $x: [a, b] \rightarrow \mathbb{R}$ be continuous and differentiable on (a, b) then $x^1 = kx$

is a simple system of ordinary differential equation in one variable.

Let $x(0) = k$ then its solution gives rise to $x(t) = x(0)e^{kt}$

$M \rightarrow TM \approx \mathbb{R}^n \rightarrow \{e_i : e_1 = (1, \dots, 0), e_n = (0, \dots, 0, 1)\}$

$\{\partial/\partial x_i : i = 1, 2, \dots, n\}$ local vector fields.

$X \in \mathbb{R}^n$ then $X = (x_1, x_2, \dots, x_n)$ where each $X_i \in \mathbb{R}$

$x^1(t) = (x^1_1(t), \dots, x^1_n(t))$ where each $x^1_i(t) = \frac{dX_i(t)}{dt}$

Reflecting this in a system of n variables and treating $\frac{d}{dt}$ as a operator which is infact linear operator then $X(t) = AX(t)$

Where A is the $n \times n$ matrix associated with the derivative operator.

The linear isomorphism in vertation can be generalised to M .

1.3 Application :

In this section to give one application form over discussion

Let the manifold be $M = \mathbb{R}^2$

For a simply connected and bounded subset D of \mathbb{R}^2 with ∂D simple closed curve then for any $\gamma: I \rightarrow \mathbb{R}^2$ as earlier such that $\gamma(I) \approx D$

Then ∂D is a simple curve in \mathbb{R}^2

If X is Vector field generated by γ in \mathbb{R}^2 then $t \rightarrow \gamma(t)$ and $X(\gamma(t))$ smoothly give rise to $X^1(\gamma(t)) = AX(\gamma(t))$

Where A is a 2×2 matrix non singular.

Which is an system of ordinary differential equation of first order setting $\gamma(t) = u$ for some point u in \mathbb{R}^2 then this parametric family $\{\gamma(t): t \in I\}$ gives rise to $\{u: u \in \mathbb{R}^2\}$

Consequently the system of ordinary differential equation, reduces to $x^1(u) = AX(u)$

References

- [1]. Benjamin and R.Milgram Journal of Differential Geometry 33(1991) 301-324 Topology of spaces.
- [2]. S.Kobagashi and Nomizu Wiley Inter science Foundation of Differential Geometry, Volume 1 and Volume 2.
- [3]. K.Fukoya Journal of Differential Geometry volume 28 (1988) 1-21 , Reimanian manifolds with bounded curvature.