



<http://www.bomsr.com>

Email:editorbomsr@gmail.com

RESEARCH ARTICLE



ON PARAMETRIC METRIC SPACES


HANDE POSUL^{1*}, SERVET KUTUKCU²

¹Department of Mathematics Faculty of Science and Arts Kilis 7 Aralik University 79000 Merkez, Kilis, Turkey

*E-mail: handeposul@kilis.edu.tr

²Department of Mathematics Faculty of Science and Arts OndokuzMayis University 55139 Kurupelit, Samsun, Turkey

E-mail: skutukcu@kilis.edu.tr



ABSTRACT

The aim of this paper is to give some basic properties in parametric metric spaces defined in literature illustrating with examples.

Keywords: Parametric metric space, open set, Hausdorffness.

AMS(2010) Subject Classification: 47H10, 54H25.

1. INTRODUCTION

In 2011, the structure of parametric metric space was introduced by Hussain et al.[1] as a new generalized metric space. After that many researchers studied this new notion ([2], [3], [4]).

Definition 1.1. [1] Let X be a nonempty set. A function $P : X \times X \times (0, \infty) \rightarrow [0, \infty)$ is said to be a parametric metric function on X , if the following conditions are satisfied; for all $x, y, z \in X$ and $t > 0$,

P1. $P(x, y, t) \geq 0$,

P2. $P(x, y, t) = 0 \Leftrightarrow x = y$,

P3. $P(x, y, t) = P(y, x, t)$,

P4. $P(x, y, t) \leq P(x, z, t) + P(z, y, t)$

Then, (X, P) is called a parametric metric space.

Example 1.1. [1] Let $X = \{f \mid f : (0, \infty) \rightarrow [0, \infty)\}$. Define $P : X \times X \times (0, \infty) \rightarrow [0, \infty)$ by

$$P(f, g, t) = |f(t) - g(t)|$$

for all $f, g \in X$ and $t > 0$. In that case, (X, P) is a parametric metric space.

Example 1.2. [1] Let $X = [0, \infty)$. Define $P : X \times X \times (0, \infty) \rightarrow [0, \infty)$ by

$$P(x, y, t) = \begin{cases} t \cdot \max\{x, y\}, & x \neq y \\ 0, & x = y \end{cases}$$

for all $x, y \in X$ and $t > 0$. In that case, (X, P) is a parametric metric space.

Definition 1.2.[1] Let (X, P) be a parametric metric space, $x \in X$ and $r > 0$. Then,

$$B_p(x, r) = \{y \in X : P(x, y, t) < r, \forall t > 0\}$$

is an open ball with center x and radius r .

Definition 1.3. [1] Let P be a parametric metric on a set X . A sequence $\{x_n\} \in X$ converges to $x \in X$ if for any $\varepsilon > 0$, there exists a $n_0 \in \mathbb{N}$ such that $P(x, x_n, t) < \varepsilon$ for all $n \geq n_0$, x is said to be the limit of $\{x_n\}$ and this is denoted by $\lim_{n \rightarrow \infty} x_n = x$ or $x_n \rightarrow x$ as $n \rightarrow \infty$.

Definition 1.4. [1] Let P be a parametric metric on a set X . A sequence $\{x_n\} \in X$ is called Cauchy sequence if for any $\varepsilon > 0$, there exists a $n_0 \in \mathbb{N}$ such that $P(x_n, x_m, t) < \varepsilon$ for all $n, m \geq n_0$.

Definition 1.5. [1] A parametric metric space (X, P) is said to be complete if every Cauchy sequence in X is convergent.

Theorem 1.1.[1] Let (X, P) be a parametric metric space, $\{x_n\}$ and $\{y_n\}$ be sequences in this space and $x, y \in X$. If $x_n \rightarrow x$ and $y_n \rightarrow y$ as $n \rightarrow \infty$ then $\lim_{n \rightarrow \infty} P(x_n, y_n, t) = P(x, y, t)$.

Proof: Since $\{x_n\}$ converges to x , for any $\varepsilon > 0$, there exists a $n_1 \in \mathbb{N}$ such that $P(x, x_n, t) < \frac{\varepsilon}{2}$ for all $n \geq n_1$. Similarly $\{y_n\}$ converges to y , for any $\varepsilon > 0$, there exists a $n_2 \in \mathbb{N}$ such that

$$P(y, y_n, t) < \frac{\varepsilon}{2} \text{ for all } n \geq n_2. \text{ Choose } n_0 = \max\{n_1, n_2\}. \text{ Then for all } n \geq n_0,$$

$$\begin{aligned} P(x_n, y_n, t) &\leq P(x_n, x, t) + P(x, y_n, t) \\ &\leq P(x_n, x, t) + P(x, y, t) + P(y, y_n, t) \\ &= \frac{\varepsilon}{2} + \frac{\varepsilon}{2} + P(x, y, t). \end{aligned}$$

From this $P(x_n, y_n, t) - P(x, y, t) < \varepsilon$. Similarly,

$$\begin{aligned} P(x, y, t) &\leq P(x, x_n, t) + P(x_n, y, t) \\ &\leq P(x, x_n, t) + P(x_n, y_n, t) + P(y_n, y, t) \\ &= \frac{\varepsilon}{2} + \frac{\varepsilon}{2} + P(x_n, y_n, t). \end{aligned}$$

From this $P(x, y, t) - P(x_n, y_n, t) < \varepsilon$. Then,

$$|P(x_n, y_n, t) - P(x, y, t)| < \varepsilon$$

is obtained. So $\lim_{n \rightarrow \infty} P(x_n, y_n, t) = P(x, y, t)$.

2. MAIN RESULTS

Definition 2.1. Let (X, P) be a parametric metric space and $F, G \subset X$. G is called an open set if for any $\varepsilon > 0$, there exists a $r > 0$ such that $B_p(x, r) \subset G$. If $X \setminus F$ is an open set, then F is called a close set.

Definition 2.1. Let (X, P) be a parametric metric space.

$$\tau = \{A \subset X : \text{for all } x \in A, \text{ there exists a } r > 0 \text{ such that } B_p(x, r) \subset A\}$$

is a topology on X .

Theorem 2.1. Let (X, P) be a parametric metric space. In this space every open ball is an open set.

Proof: Consider the open ball $B_p(x, r)$. Then $y \in B_p(x, r) \Rightarrow P(x, y, t) < r$. Assume that $r' = r - P(x, y, t)$. Take into consideration the ball $B_p(y, r')$. We claim that $B_p(y, r') \subset B_p(x, r)$.

Let $z \in B_p(y, r')$. Then, $P(y, z, t) < r'$. For this reason,

$$\begin{aligned} P(x, z, t) &\leq P(x, y, t) + P(y, z, t) \\ &< P(x, y, t) + r' \\ &= P(x, y, t) + r - P(x, y, t) \\ &= r. \end{aligned}$$

Then, $z \in B_p(x, r)$. So, the proof is completed.

Theorem 2.2. Every parametric metric space is a Hausdorff space.

Proof: Let (X, P) be a parametric metric space and $x, y \in X$, $x \neq y$. Consider $P(x, y, t) = r$. Take

into consideration the open sets $U = B_p(x, \frac{r}{2})$ and $V = B_p(y, \frac{r}{2})$. It is clear that $x \in U$ and $y \in V$.

We claim that $U \cap V = \emptyset$. Assume that $U \cap V \neq \emptyset$. Then there exists a $z \in X$ such that

$z \in U \cap V$. Then $z \in U \Rightarrow P(x, z, t) < \frac{r}{2}$ and $z \in V \Rightarrow P(y, z, t) < \frac{r}{2}$. From P4,

$$\begin{aligned} P(x, y, t) &\leq P(x, z, t) + P(z, y, t) \\ &= P(x, z, t) + P(y, z, t) \\ &= \frac{r}{2} + \frac{r}{2} \\ &= r. \end{aligned}$$

Since $P(x, y, t) = r$, $r < r$ is obtained. But this is a contradiction. Then $U \cap V = \emptyset$. The proof is completed.

Theorem 2.3. Let (X, P) be a parametric metric space. In this space every convergent sequence has a unique limit point.

Proof: Let $\{x_n\} \subset X$ be a sequence and $x, y \in X$, $x \neq y$. Suppose that $\{x_n\}$ converges to both x

and y . Since $\{x_n\}$ converges to x , for any $\varepsilon > 0$, there exists a $n_1 \in \mathbb{N}$ such that $P(x, x_n, t) < \frac{\varepsilon}{2}$

for all $n \geq n_1$. Similarly $\{x_n\}$ converges to y , for any $\varepsilon > 0$, there exists a $n_2 \in \mathbb{N}$ such that

$P(x, x_n, t) < \frac{\varepsilon}{2}$ for all $n \geq n_2$. Choose $n_0 = \max\{n_1, n_2\}$. Then for all $n \geq n_0$,

$$\begin{aligned} P(x, y, t) &\leq P(x, x_n, t) + P(x_n, y, t) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon. \end{aligned}$$

Since ε is arbitrary, $P(x, y, t) = 0 \Leftrightarrow x = y$. Then every convergent sequence in this space has a unique limit point.

Theorem 2.4. Let (X, P) be a parametric metric space. In this space every convergent sequence is a Cauchy sequence.

Proof: Let $\{x_n\} \subset X$ be a convergent sequence. Then there exists a $x \in X$ such that $\lim_{n \rightarrow \infty} x_n = x$.

So for any $\varepsilon > 0$, there exist $n_1, n_2 \in \mathbb{N}$ such that $P(x, x_n, t) < \frac{\varepsilon}{2}$ for all $n \geq n_1$ and

$P(x, x_n, t) < \frac{\varepsilon}{2}$ for all $m \geq n_2$. Choose $n_0 = \max\{n_1, n_2\}$. Then for all $m, n \geq n_0$,

$$\begin{aligned} P(x_n, x_m, t) &\leq P(x_n, x, t) + P(x, x_m, t) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon. \end{aligned}$$

Then $\{x_n\}$ is a Cauchy sequence.

REFERENCES

- [1]. Hussain N., Khaleghizadeh S., Salimi P., Afrah A. N. A., 2014, A new approach to fixed point results in triangular intuitionistic fuzzy metric spaces, *Abstract and Applied Analysis*, doi:10.1155/2014/690139.
- [2]. Hussain N., Salimi P., Salimi P., Parvaneh V., 2015, Fixed point results for various contractions in parametric and fuzzy b-metric spaces, *J. Nonlinear Sci. Appl.*, 8(5), 719-739.
- [3]. Rashmi J., Daheriya R. D., Ughade M., 2016, Fixed point, coincidence point and common fixed point theorems under various expansive conditions in parametric metric spaces and parametric b-metric spaces, *Gazi University Journal of Science*, 29(1), 95-107.
- [4]. Likhitker M., Daheriya R. D., Picard sequence and fixed point results on parametric b-metric spaces, *International Journal of Mathematics*, 4(1), 1220-1229.